

Anti-Oxidative Water Improves Diabetes

S. SHIRAHATA, T. NISHIMURA, S. KABAYAMA*, D. AKI, K. TERUYA,
K. OTSUBO*, S. MORISAWA*, Y. ISHII**, Z. GADEK*** and Y. KATAKURA

*Graduate School of Genetic Resources Technology, Kyushu University, Fukuoka 812-8581, Japan; *Nihon Trim Co. Ltd., 1-8-34 Oyodonaka, Kita-ku, Osaka 531-0076, Japan; **Hita Tenryosui Co. Ltd., Nakanoshima-machi, Hita 877-0074, Japan; ***Centre for Holistic Medicine and Naturopathy, 57392 Nordenau, Germany*

Abstract. Electrolyzed reduced water and natural waters such as Hita Tenryosui water in Japan, Nordenau water in Germany and Tracote water in Mexico, which are known to improve various diseases, were all anti-oxidative waters which could scavenge intracellular reactive oxygen species. The anti-oxidative waters stimulated not only the glucose uptake of rat myotube L6 cells, but also the secretion of insulin from a pancreatic beta cell line HIT-T15. The anti-oxidative waters improved the damage in the sugar tolerance test of type 2 diabetes model mice (*db/db* mice). A clinical investigation demonstrated that Nordenau water could significantly improve the diabetes mellitus.

1. Introduction

Reactive oxygen species (ROS) such as $^1\text{O}_2$, O_2^- , H_2O_2 and $\cdot\text{OH}$ are known to cause various diseases including diabetes mellitus. Electrolyzed reduced water (ERW) produced near cathode during electrolysis of water scavenged ROS and protected DNA from oxidative damage *in vitro* (1). Hayashi proposed a water regulation theory that hydrogen-rich water can scavenge ROS (2) and reported that the daily intake of ERW improved diabetes mellitus (3). ERW and natural reduced water such as Hita Tenryosui water stimulated glucose uptake of rat L6 myotubes (4). There are several natural waters drawn from deep underground such as Nordenau water in Germany and Tracote water in Mexico as well as Hita Tenryosui water in Japan, which are called as miracle water because of curing power against various diseases. Here we demonstrate that ERW and the natural waters are all anti-oxidative water and scavenge ROS in cultured cells. The anti-oxidative waters stimulated not only the glucose uptake into insulin-responsive muscle cells but also the insulin secretion from beta cells of pancreas. The anti-oxidative water improved the damage in the sugar-tolerance test of diabetes model mice and also improved the symptoms of human diabetes.

2. Materials and Methods

Waters. ERW was prepared by electrolysis of activated charcoal treated tap water or ultra pure water containing 0.01% NaCl using an electrolyzing device (TI-8000, Nihon Trim Co., Ltd., Osaka). The pH and ORP of ERW were 10.5 to 11.5 and -650 to -850 mV, respectively. ERW was used after neutralized with HCl. Hita Tenryosui water was supplied by Hita Tenryosui Co. Ltd., Hita, Japan. Nordenau water was given by Mr. Theo Tommes, Nordenau, Germany. Tracote water was supplied by Nihon TV Co. Ltd. Mineral waters A and B were those commercially sold in Japan.



Measurement of intracellular ROS scavenging effects. L6 cells derived from rat skeletal muscle were differentiated into insulin responsive myotubes by cultivation in 2% FBS/DME medium for 7 days (4). Intracellular redox state of L6 cells cultured in medium prepared with each waters was measured after 72 hour cultivation by a confocal laser microscopy using a fluorescent dye, 2',7'- dichlorofluorescein diacetate (DCFH-DA) as a probe (5).

Measurement of insulin secretion from beta cells of pancreas. HIT-T15 cell line derived from Syrian hamster pancreatic beta cells were cultured in Ham' F12 medium supplemented with 10% dialyzed horse serum and 2.5% FBS. The cells (1×10^4 cells/ml) were inoculated into a 6 well microplate. After 3 days, the medium was changed to medium containing each water and the cultivation was done in the presence of 10 mM glucose for 72 hours. Insulin was measured by a insulin EIA kit (Amarsham Pharmacia).

Animal experiments using type 2 diabetes model mice. Type 2 diabetes model mice (C57BL/KsJ-Db⁺/Db⁺) were obtained from Nihon Crea Co. Ltd. The mice, which have a defect in the leptin receptor gene, cannot control appetite and suffer from diabetes via obesity. Blood sugar level was measured using a dexistar Z sensor equipped with a enzyme electrode (Bayer-Sankyo, Tokyo). Sugar tolerance test was performed by intraperitoneal injection of glucose solution (1 mg glucose in physiological salt solution/30 g weight). Blood sugar lever was determined at 0, 30 and 60 minutes after the injection (6).

Clinical analysis on the effect of Nordenau water against diabetes. Changes in the relevant test parameters of 139 diabetes patients were examined. The average age of the test persons (72 female, 67 male) was 68.4 years; the average duration of their stay in Nordenau was 4 days for group 1 (79 patients = 56.8%), 5 or more days for group 2 (60 patients = 43.2%) taking tow tunnel walks of 30 minutes each and consuming 2 litres of Nordenau water daily. The patients were being looked after by us diagnostically within the scope of what is known as 'course control' and they gave us permission to use their diagnostic parameters for scientific purposes, provided we guaranteed their anonymity. We tested the diabetes-relevant test parameters (blood sugar, HbA1c, chol., trigl., HDL and LDL) twice under the same conditions i.e. on their arrival and on their departure at the same time of day. During the period of observation neither their regular medication nor their diet instructions was altered. The tested persons were particularly reminded to continue the medication prescribed by their general practitioners, to stick to their individual diet plans and not to alter any of their behavioural patterns. The statistical interpretation of the data contains: (a) The descriptive statistics of the whole group and of both subgroups. (b) A pairing-directed T-Test. (c) A proportional evaluation of the entire group and of both subgroups. (d) Variance- and co-variance analyses. The allowance for error probability was set at 5%.




Figure 1. Improvement of sugar tolerance test of type 2 diabetes model mice by anti-oxidative water. A, tap water; B, ERW containing 10 ppm H_2PtCl_6 ; C, Hita Tenryosui water; D, Nordenau water.

3. Results and Discussion

3.1. ROS SCAVENGING EFFECTS OF ERW AND NATURAL WATERS

Differentiated L6 myotubes exhibited low level of ROS stress and were sensitive for ROS scavengers. Ascorbic acid (AsA) exhibited maximum 22% intracellular ROS scavenging effect (22%) at 60 μM , however, higher and lower concentrations of AsA exhibited the decreased scavenging effects and more than 1 mM AsA exhibited no scavenging effects, suggesting that AsA acts as both a reductant and an oxidant. ERW produced from tap water scavenged 25% of intracellular ROS. Hita Tenryosui water, Nordenau water and Tracote water scavenged 20%, 24% and 24% intracellular ROS, respectively. However, commercial mineral waters A and B shifted intracellular redox state to 10% and 6% oxidized state. Tap water containing hypochloric acid shifted intracellular redox state to 25% oxidized state. These results demonstrated that ERW and natural waters, which is known to improve various diseases, are all anti-oxidative waters.

3.2. STIMULATION OF THE SECRETION OF INSULIN FROM PANCREATIC BETA CELLS BY ANTI-OXIDATIVE WATER

The effect of waters on the secretion of insulin by HIT-T15 cells were examined. In the medium containing ultra pure water, the control cells secreted 0.03 ng/ml of insulin. ERW and Nordenau water-treated cells secreted 3.3 and 2.9 times higher concentration of insulin than control cells. Hita Tenryosui water also stimulated the secretion of insulin about 2.2 times compared to control, however, the commercial mineral water did not stimulate the secretion of insulin but decreased the secretion to 0.73 times of control. Since ROS suppressed the secretion of insulin, the anti-oxidative waters may increase the insulin secretion by decreasing the oxidative stress in the pancreatic cells.

3.3. IMPROVEMENT OF DAMAGE IN SUGAR TOLERANCE TEST OF TYPE 2 DIABETES MODEL MICE BY ANTI-OXIDATIVE WATER

As shown in Figure 1, the type 2 diabetes model mice administered tap water exhibited the damaged result in the sugar tolerance test. Whereas, the mice administered ERW containing 10 ppm of H_2PtCl_6 , Hita Tenryosui water or Nordenau water exhibited the




Figure 2. Improvement of human diabetes by drinking Nordenau water. Diabetes patients drunk 2 liter of Nordenau water every day. A, blood sugar average value – group 2; B, HbA1c average value – group 2; C, cholesterol average value – group 1. Group 1, drinking for 4 days. Group 2, drinking for 5 or more days. Cont., values on the arrival; Test, values on the departure.

improved result in the sugar tolerance test. Administration of ERW or H_2PtCl_6 only did not exhibit significant improvement (data not shown). Since ERW was prepared using ultra pure water, H_2PtCl_6 was expected to work as a carrier of active hydrogen in ERW. Since natural reduced waters such as Hita Tenryosui water and Nordenau water contains various kinds of minerals, they may exhibit more prominent effects *in vivo* than ERW.

3.4. EFFECTS OF NORDENAU WATER ON DIABETES PATIENTS

The descriptive statistics of the group as a whole shows a downward trend of the average blood sugar, HbA1c, chol., trigl., and LDL. The HDL average, however, shows an upward trend. By assuming a unilateral and hypothetical error probability alpha of 5%, the pairing test showed a significant improvement merely as regards the counts of blood sugar and HbA1c in subgroup 2 (average duration of stay 5 days and more). Cholesterol has improved significant in the group 1 only. In subgroup I (average duration of stay 4 days) beside cholesterol practically no changes of diabetes-relevant parameters have been observed.

References

1. Shirahata, S. et al, *Biochem. Biophys. Res. Commun.*, **234**, 269–274 (1997).
2. Hayashi, H., *Explore*, **6**, 28–31 (1995).
3. Hayashi, H. and Kawamura, M., *Animal Cell Technology: Basic & Applied Aspects*, **12**, in press.
4. Oda et al., *Animal Cell Technology: Products from Cells, Cells as Products*, A. Bernard et al. (eds.), pp. 425–427, Kluwer Academic Publishers (1999).
5. Charles X. Xie et al., *Brain Res.*, **743**, 271–277 (1996).
6. Takahashi, M. et al., *Proc. Natl. Acad. Sci. USA*, **94**, 11969–11973 (1997).